

Experimental review of pentaquarks and tetraquarks

Jibo HE
on behalf of the LHCb collaboration, including results from
the Babar, Belle, BESIII, CMS, D0, CDF collaborations

University of Chinese Academy of Sciences (UCAS), Beijing

FPCP 2016 @ Caltech, 09/06/2016

Introduction

- Quark model proposed by Gell-Mann and Zweig in 1964
- Tetraquarks and pentaquarks are not a priori excluded...

Volume 8, number 3

PHYSICS LETTERS

1 February 1964

A SCHEMATIC MODEL OF BARYONS AND MESONS *

M. GELL-MANN

California Institute of Technology, Pasadena, California

Received 4 January 1964

We then refer to the members $u^{\frac{1}{3}}$, $d^{-\frac{1}{3}}$, and $s^{-\frac{1}{3}}$ of the triplet as "quarks" 6) q and the members of the anti-triplet as anti-quarks \bar{q} . Baryons can now be constructed from quarks by using the combinations (qqq) , $(qqq\bar{q}\bar{q})$, etc., while mesons are made out of $(q\bar{q})$, $(q\bar{q}\bar{q}\bar{q})$, etc. It is assuming that the lowest

Experimental status


- XYZ keep appearing, continuous efforts to understand their nature

State | Experiment

$X(3872)$	Belle, CDF, D0, Babar, LHCb, BESIII, CMS
$X(3915)$	Belle, Babar
$X(3940)$	Belle
$G(3900)$	Babar, Belle
$Y(4008)$	Belle
$X(4140)$	CDF, LHCb, CMS, D0, Babar, Belle, BESIII
$X(4160)$	Belle
$Y(4260)$	Babar, CLEO, Belle
$X(4274)$	CDF, CMS, Babar, BESIII, LHCb
$X(4350)$	Belle
$Y(4360)$	Babar, Belle
$X(4630)$	Belle
$Y(4660)$	Belle
$Z_c^+(3900)$	BESIII, Belle, CLEO
$Z_c^+(4020)$	BESIII
$Z_c^+(4050)$	Belle, Babar
$Z^+(4200)$	Belle
$Z_2^+(4250)$	Belle, Babar
$Z^+(4430)$	Belle, Babar, LHCb
$Y_b(10890)$	Belle
$Z_b^+(10610)$	Belle
$Z_b^0(10610)$	Belle
$Z_b^+(10650)$	Belle
$P_c(4450)^+$	LHCb
$P_c(4380)^+$	LHCb

Not a complete list...

[S.L. Olsen, Front. Phys. 10 (2015) 101401]



- Please see Dr. Estia Eichten's talk for theoretical discussions






The $X(3872)$

$(c\bar{c}[u\bar{u} + d\bar{d}])$

[Belle, PRL 91 (2003) 262001] [CDF, PRL 93 (2004) 072001] [D0, PRL 93 (2004) 162002]

- $X(3872)$ discovered by Belle with $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$ in 2003, quickly confirmed by CDF and D0 in $p\bar{p}$ collisions





- Nature of $X(3872)$ still unclear

- ▶ Molecular state, $D\bar{D}^{0*}$ threshold ($J^{PC} = 0^{-+}, 1^{++}$)
- ▶ Tetraquark (1^{++})
- ▶ $\chi_{c1}(2^3P_1)$ (1^{++}) or $\eta_{c2}(1^1D_2)$ (2^{-+})

Determining J^{PC} of $X(3872)$

[LHCb, PRL 110 (2013) 222001]


- Thanks to CDF, Belle and Babar, J^{PC} of $X(3872)$ narrowed to 1^{++} or 2^{-+}
- Full 5D angular analysis by LHCb, with ~ 310 $B^+ \rightarrow X(3872)K^+$ in 2011 data (1 fb^{-1}), 2^{-+} rejected by 8σ



- All assumed the lowest orbital angular momentum L_{\min} in $X(3872) \rightarrow J/\psi \rho$


Determining J^{PC} of $X(3872)$, w/o assuming L_{\min}

- LHCb updated analysis without assuming L_{\min} , w/ 3 fb^{-1} data



J^{PC}	B_{LS} Any L value	B_{LS} Minimal L value
0^{++}	B_{11}	B_{11}
0^{++}	B_{00}, B_{22}	B_{00}
1^{-+}	$B_{10}, B_{11}, B_{12}, B_{32}$	B_{10}, B_{11}, B_{12}
1^{++}	B_{01}, B_{21}, B_{22}	B_{01}
2^{-+}	$B_{11}, B_{12}, B_{31}, B_{32}$	B_{11}, B_{12}
2^{++}	$B_{02}, B_{20}, B_{21}, B_{22}, B_{42}$	B_{02}
3^{-+}	$B_{12}, B_{30}, B_{31}, B_{32}, B_{52}$	B_{12}
3^{++}	$B_{21}, B_{22}, B_{41}, B_{42}$	B_{21}, B_{22}
4^{-+}	$B_{31}, B_{32}, B_{51}, B_{52}$	B_{31}, B_{32}
4^{++}	$B_{22}, B_{40}, B_{41}, B_{42}, B_{62}$	B_{22}

[LHCb, PRD 92 (2015) 011102(R)]




- $J^{PC} = 1^{++}$ confirmed, D-wave fraction <4% at 95% C.L.

$X(3872) \rightarrow J/\psi \gamma, \psi(2S)\gamma$

[LHCb, NPB 886 (2014) 665]




- $R_{\psi\gamma} \equiv \frac{\mathcal{B}(X \rightarrow \psi(2S)\gamma)}{\mathcal{B}(X \rightarrow J/\psi\gamma)}$ expected different for $c\bar{c}$ or $D\bar{D}^{*0}$ molecule
- Following Babar, Belle, LHCb also studied $X(3872) \rightarrow J/\psi\gamma, \psi(2S)\gamma$ with 3 fb^{-1} of data
- Evidence of $X(3872) \rightarrow \psi(2S)\gamma$ found, 4.4σ .
Pure $D\bar{D}^{*0}$ molecule disfavored




$X(3872)$ hadroproduction

- Following CDF, LHCb, CMS also studied
 $X(3872)$ production at 7 TeV

[CMS, JHEP 04 (2013) 154]



[P. Artoisenet and E. Braaten, PRD 81 (2010) 114018]




The $X(4140)$ and $X(4274)$

$(c\bar{c}ss\bar{s})$

[CDF PRL 102 (2009) 242002, arXiv: 1101.6058] [LHCb, PRD 85 (2012) 091103(R)]
 [D0, PRD 89 (2014) 012004] [CMS, PLB 734 (2014) 261]





- Seen by CDF in $B^+ \rightarrow K^+ J/\psi \phi$, not by LHCb... All 1D fits



$B^+ \rightarrow J/\psi \phi K^+$ in LHCb full data set

New

[LHCb-Paper-2016-018, in preparation]






Kaon excitations

Are these reflections of
interfering $K^* \rightarrow \phi K^-$?
Proper amplitude analysis
necessary to check

Amplitude fit including 4 exotic X

New

[LHCb-Paper-2016-018, in preparation]




- We considered adding possible exotic $X \rightarrow J/\psi \phi$ and $Z^+ \rightarrow J/\psi K^+$ states as well as removing insignificant or implausible ($\Gamma > 1000$ or < 100 MeV) conventional $K^{*+} \rightarrow \phi K^+$ states leading us to a default model
- Only X states give very significant improvements in fit qualities over the models with K^* s alone
- The default fit model is shown here.



The $X(4140)$ and $X(4274)$ properties

New

[LHCb-Paper-2016-018, in preparation]



LHCb preliminary


Contri- butio n	sign. or Ref.	M_0 MeV	Γ_0 MeV	Fit results
All $X(1^+)$				$16 \pm 3 \begin{array}{l} +6 \\ -2 \end{array}$
$X(4140)$	8.4σ	$4146.5 \pm 4.5 \begin{array}{l} +4.6 \\ -2.8 \end{array}$	$83 \pm 21 \begin{array}{l} +21 \\ -14 \end{array}$	$13 \pm 3.2 \begin{array}{l} +4.8 \\ -2.0 \end{array}$
ave.		4146.9 ± 2.3	17.8 ± 6.8	
$X(4274)$	6.0σ	$4273.3 \pm 8.3 \begin{array}{l} +17.2 \\ -3.6 \end{array}$	$56 \pm 11 \begin{array}{l} +8 \\ -11 \end{array}$	$7.1 \pm 2.5 \begin{array}{l} +3.5 \\ -2.4 \end{array}$
CDF		$4274.4 \begin{array}{l} +8.4 \\ -6.7 \end{array} \pm 1.9$	$32 \begin{array}{l} +22 \\ -15 \end{array} \pm 8$	
CMS		$4313.8 \pm 5.3 \pm 7.3$	$38 \begin{array}{l} +30 \\ -15 \end{array} \pm 16$	

- $X(4140)$, significance 8.4σ
 - ▶ Mass consistent with previous measurements, width larger
 - ▶ $J^{PC} = 1^{++}$ determined at 5.7σ
- $X(4274)$, significance 6.0σ
 - ▶ Mass/width consistent with the CDF unpublished results
 - ▶ $J^{PC} = 1^{++}$ determined at 5.8σ

Observation of two new X states

New

[LHCb-Paper-2016-018, in preparation]



LHCb preliminary




Contri- butio n	sign. .	M_0 MeV	Γ_0 MeV	Fit results F.F. %
All $X(0^+)$				$28 \pm 5^{+7}_{-7}$
NR $J/\psi\phi$	6.4σ			$46 \pm 11^{+11}_{-21}$
$X(4500)$	6.1σ	$4506 \pm 11^{+12}_{-15}$	$92 \pm 21^{+21}_{-20}$	$6.6 \pm 2.4^{+3.5}_{-2.3}$
$X(4700)$	5.6σ	$4704 \pm 10^{+14}_{-24}$	$120 \pm 31^{+42}_{-33}$	$12 \pm 5^{+9}_{-5}$

- Significant structures at higher masses, best described by two new 0^{++} resonances $X(4500)$, $X(4700)$
 - Significance of 6.1σ , 5.6σ
 - $J^{PC} = 0^{++}$ determined at 4.0σ , 4.5σ

Hadroproduction of $X(4140)$

[D0, PRL 115 (2015) 231001]

- First evidence of prompt production of $X(4140)$, 4.7σ
- Fraction of $X(4140)$ from b -decays $f_b = 0.39 \pm 0.07 \pm 0.10$




The $Z(4430)^-$

($c\bar{c}d\bar{u}$)

[Belle, PRL 100 (2008) 142001] [BaBar, PRD 79 (2009) 112001]



- Seen by Belle in $B^0 \rightarrow K^+ \psi(2S)\pi^-$, with 605 fb^{-1} @ $\Upsilon(4S)$
 - $M = 4433 \pm 4 \pm 2 \text{ MeV}$, $\Gamma = 45^{+18+30}_{-13-13} \text{ MeV}$
- Not confirmed by Babar with 413 fb^{-1} @ $\Upsilon(4S)$



The $Z(4430)^-$, further studies by Belle

[Belle, PRD 80 (2009) 031104, PRD 88 (2013) 074026]




- Dalitz analysis (2D) in 2009 (605 fb^{-1})
 - ▶ $M = 4443^{+15+19}_{-12-13} \text{ MeV}$, $\Gamma = 107^{+86+74}_{-43-56} \text{ MeV}$
- Full 4D angular analysis in 2013 (711 fb^{-1}), $N_{\text{sig}} \sim 2k$
 - ▶ $M = 4485^{+22+28}_{-22-11} \text{ MeV}$, $\Gamma = 200^{+41+26}_{-46-35} \text{ MeV}$
 - ▶ $J^P = 1^+$ favored by $> 3.4\sigma$



The $Z(4430)^-$, further studies at LHCb

[LHCb, PRL 112 (2014) 222002, PRD 92 (2015) 112009]


- LHCb did both full 4D angular analysis and model-independent analysis, with 10 times more signal $N_{\text{sig}} \sim 25k$
 - $M = 4475 \pm 7^{+15}_{-25} \text{ MeV}$
 - $\Gamma = 172 \pm 13^{+37}_{-34} \text{ MeV}$
 - $J^P = 1^+$ established
- Argand plot shows clear resonance feature



The Z_c family

$$(c\bar{c}u\bar{d}[u\bar{u} + d\bar{d}])$$


- Several Z_c states found by BESIII, Belle, CLEO
- $Z_c(3900)$ & $Z_c(3885)$ the same? $Z_c(4020)$ & $Z_c(4025)$?




$e^+e^- \rightarrow \pi^-\pi^+J/\psi$

$Z_c(3885)^+$

ST: PRL 112 (2014) 022001
DT: PRD 92 (2015) 092006




$e^+e^- \rightarrow \pi^-(D\bar{D}^*)^+$




$e^+e^- \rightarrow \pi^0\pi^0J/\psi$

$Z_c(3885)^0$

PRL 115 (2015) 222002




$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$




$e^+e^- \rightarrow \pi^-\pi^+h_c$

$Z_c(4025)^+$

PRL 112 (2014) 132001




$e^+e^- \rightarrow \pi^-(D^*\bar{D}^*)^+$



$e^+e^- \rightarrow \pi^0\pi^0h_c$

$Z_c(4025)^0$

PRL 115 (2015) 182002



$e^+e^- \rightarrow \pi^0(D^*\bar{D}^*)^0$

The Z_c family (cont.)

- $Z_c(3900)$ & $Z_c(3885)$ the same? $Z_c(4020)$ & $Z_c(4025)$?
 - ▶ Masses and widths consistent with each other within 2σ
 - ▶ Seem to favor the same J^P




Z_c	Mass (MeV/c ²)	Width (MeV/c ²)	Decay mode	J^P
$Z_c(3900)^+$	$3899.0 \pm 3.6 \pm 4.9$	$46 \pm 10 \pm 20$	$\pi^+ J/\psi$	1^+
$Z_c(3900)^0$	$3894.8 \pm 2.3 \pm 2.7$	$29.6 \pm 8.2 \pm 8.2$	$\pi^0 J/\psi$	
$Z_c(3885)^+$	$3882.3 \pm 1.1 \pm 1.9$	$26.5 \pm 1.7 \pm 2.3$	$(D\bar{D}^*)^+$	1^+
$Z_c(3885)^0$	$3885.7^{+4.3}_{-5.7} \pm 8.4$	$35^{+11}_{-12} \pm 15$	$(D\bar{D}^*)^0$	
$Z_c(4020)^+$	$4022.9 \pm 0.8 \pm 2.7$	$7.9 \pm 2.7 \pm 2.6$	$\pi^+ h_c$	
$Z_c(4020)^0$	$4023.8 \pm 2.2 \pm 3.8$	7.9 (fixed)	$\pi^0 h_c$	
$Z_c(4025)^+$	$4026.3 \pm 2.6 \pm 3.7$	$24.8 \pm 5.6 \pm 7.7$	$(D^*\bar{D}^*)^+$	
$Z_c(4025)^0$	$4025.5^{+2.0}_{-4.7} \pm 3.1$	$23.0 \pm 6.0 \pm 1.0$	$(D^*\bar{D}^*)^0$	

The $X(5568)^+$ by D0

$(\bar{b} s u d)$

[D0, arXiv: 1602.07588]



- $X(5568)^+ \rightarrow B_s^0 \pi^+$ observed by D0 with 5.1σ
 - ▶ $M = 5567.8 \pm 2.9^{+0.9}_{-1.9}$ MeV
 - ▶ $\Gamma = 21.9 \pm 6.4^{+5.0}_{-2.5}$ MeV
 - ▶ $\rho_X^{\text{D0}} \equiv \frac{\sigma(X)\mathcal{B}(X \rightarrow B_s^0 \pi^+)}{\sigma(B_s^0)}|_{\text{D0Acc}}$
 $= (8.6 \pm 1.9 \pm 1.4)\%$
for $p_T(B_s^0) > 10$ GeV



The $X(5568)^+$, not confirmed by LHCb



[LHCb-CONF-2016-004]

- Using both $B_s^0 \rightarrow D_s^- \pi^+$ and $B_s^0 \rightarrow J/\psi \phi$, cut based selection for clean B_s^0 samples, $N(B_s^0)$ 10 times more than D0



$$p_T(B_s^0) > 10 \text{ GeV}/c$$


No significant signal



The $X(5568)^+$, upper limits on production rate


[LHCb-CONF-2016-004]

- Upper limits on $X(5568)^+$ production rate at LHC
 - $\rho_X^{\text{LHCb}}(B_s^0 p_T > 5 \text{ GeV}) < 0.009(0.010) @ 90 (95)\% \text{ C.L.}$
 - $\rho_X^{\text{LHCb}}(B_s^0 p_T > 10 \text{ GeV}) < 0.016(0.018) @ 90 (95)\% \text{ C.L.}$






$\Lambda_b \rightarrow J/\psi p K^+$, unexpected structure in $m_{J/\psi p}$

[LHCb, PRL 115 (2015) 072001]



$\Lambda(1520)$ and other Λ^* 's $\rightarrow p K^-$



Exotic pentaquark



- Unexpected, narrow peak in $m_{J/\psi p}$


Is it a reflection of
interfering Λ^* 's $\rightarrow p K^-$?
Proper amplitude analysis
necessary to check






Observation of $P_c(4450)^+$ and $P_c(4380)^+$ ($c\bar{c}uud$)

[LHCb, PRL 115 (2015) 072001]

w/o P_c 's



w/ P_c 's




Observation of $P_c(4450)$ and $P_c(4380)$ (cont.)

[LHCb, PRL 115 (2015) 072001, Chin. Phys. C 40 (2016) 011001]

- Two new states needed to describe the data

	$P_c(4380)^+$	$P_c(4450)^+$
J^P	$\frac{3}{2}^-$	$\frac{5}{2}^+$
Mass [MeV/ c^2]	$4380 \pm 8 \pm 29$	$4449.8 \pm 1.7 \pm 2.5$
Width [MeV]	$205 \pm 18 \pm 86$	$39 \pm 5 \pm 19$
Significance	9σ	12σ

- Argand plots



- Branching fractions also measured

$$\mathcal{B}(\Lambda_b^0 \rightarrow P_c^+(4380) K^-) \mathcal{B}(P_c^+ \rightarrow J/\psi p) = (2.56 \pm 0.22 \pm 1.28^{+0.46}_{-0.36}) \times 10^{-5}$$


$$\mathcal{B}(\Lambda_b^0 \rightarrow P_c^+(4450) K^-) \mathcal{B}(P_c^+ \rightarrow J/\psi p) = (1.25 \pm 0.15 \pm 0.33^{+0.22}_{-0.18}) \times 10^{-5}$$



Pentaquark, model-independent analysis

[LHCb, arXiv:1604.05708]



- Model-independent approach: not assume anything about Λ^* , Σ^* or NR compositions, spin, masses, widths or mass-shape
- Only restrict maximal spin of allowed Λ^* at given m_{Kp}
- Null hypothesis (H_0) rejected at 9σ



Cabbibo suppressed vs favored Λ_b decay





[LHCb, PRL 115 (2015) 072001] [LHCb-Paper-2016-015, in preparation]

- Cabbibo suppressed $\Lambda_b \rightarrow J/\psi p\pi^+$, signal yields 10 times smaller than $\Lambda_b \rightarrow J/\psi pK^+$
- Relatively higher background...



Cabbibo suppressed $\Lambda_b \rightarrow J/\psi p\pi^+$

[LHCb-Paper-2016-015, in preparation]



- No obvious structure in $m_{J/\psi p}$

Statistics is low.

Proper amplitude analysis necessary to check for consistency with Cabibbo favored

$$\Lambda_b^0 \rightarrow J/\psi K^-$$






Pentaquarks in $\Lambda_b \rightarrow J/\psi p\pi^+$ decay

New

[LHCb-Paper-2016-015, in preparation]

- Not enough statistics to do open-ended analysis of $J/\psi p$, $J/\psi \pi^-$ contributions, check data for previously observed states
- Combined significance of $P_c(4380)^+$, $P_c(4450)^+$, $Z_c(4200)^-$ together, 3.1σ

LHCb preliminary



Summary

- Many charmonium-like and bottomonium-like XYZ states found in the past decade, their nature remains unclear
 - ▶ Pentaquarks, tetraquarks; molecule; $c\bar{c}/b\bar{b}$?
- Great progress made to help understand their nature
 - ▶ J^{PC} , $X(3872)$, $Z(4430)^-$, $X(4140)$, ...
 - ▶ New decay modes, $X(3872) \rightarrow \psi(2S)\gamma$, ...
 - ▶ Hadroproduction, $X(3872)$, $X(4140)$
- Tensions between results of different experiments resolved, e.g., $Z(4430)^-$, $X(4140)$; new tension appears, $X(5568)^+$
- Your suggestions on high-priority analysis are always welcome